

MEETING ABSTRACT

Open Access

Assessment of the quality of brain regions and neuroimaging metrics as biomarkers of Alzheimer's Disease

Tânia Faria Vaz^{1,2*}, Filipa Lucena^{1,2}, Joana Pé-Leve¹, André Santos Ribeiro³, Luís Lacerda⁴, Nuno da Silva⁵, David Nutt³, John McGonigle³, Hugo Alexandre Ferreira¹

From PSMR 2015: 4th Conference on PET/MR and SPECT/MR
La Biodola, Isola d'Elba, Italy. 17-21 May 2015

¹Institute of Biophysics and Biomedical Engineering, Faculty of Sciences, University of Lisbon, Portugal

Alzheimer Disease (AD) is characterized by progressive cognitive decline and dementia. Earlier diagnosis and classification of different stages of the disease are currently the main challenges and can be assessed by neuroimaging. With this work we aim to evaluate the quality of brain regions and neuroimaging metrics as biomarkers of AD. Multimodal Imaging Brain Connectivity Analysis (MIBCA) toolbox functionalities were used to study AD by T1weighted, Diffusion Tensor Imaging and 18FAV45 PET, with data obtained from the AD Neuroimaging Initiative database, specifically 12 healthy controls (CTRL) and 33 patients with early mild cognitive impairment (EMCI), late MCI (LMCI) and AD (11 patients/group). The metrics evaluated were gray-matter volume (GMV), cortical thickness (CThk), mean diffusivity (MD), fractional anisotropy (FA), fiber count (FiberConn), node degree (Deg), cluster coefficient (ClusC) and relative standard-uptake-values (rSUV). Receiver Operating Characteristic (ROC) curves were used to evaluate and compare the diagnostic accuracy of the most significant metrics and brain regions and expressed as area under the curve (AUC). Comparisons were performed between groups. The RH-Accumbens/Deg demonstrated the highest AUC when differentiating between CTRL-EMCI (82%), whether rSUV presented it in several brain regions when distinguishing CTRL-LMCI (99%). Regarding CTRL-AD, highest AUC were found with LH-STG/FiberConn and RH-FP/FiberConn (~100%). A larger number of neuroimaging metrics related with cortical atrophy with AUC>70% was found in CTRL-AD in both hemispheres, while in earlier stages, cortical metrics showed in more confined areas of the temporal region and mainly in LH, indicating an increasing of the spread of cortical atrophy that is characteristic of disease progression. In CTRL-EMCI several brain regions and neuroimaging metrics presented AUC>70% with a worst result in later stages suggesting these indicators as biomarkers for an earlier stage of MCI, although further research is necessary.

Authors' details

¹Institute of Biophysics and Biomedical Engineering, Faculty of Sciences, University of Lisbon, Portugal. ²Lisbon School of Health Technology, Polytechnic Institute of Lisbon, Portugal. ³Centre for Neuropsychopharmacology, Division of Brain Sciences, Department of Medicine, Imperial College London, UK. ⁴Centre for Neuroimaging Sciences, Institute of

Psychiatry, King's College London, UK. ⁵Institute of Neuroscience and Medicine 4, Forschungszentrum Jülich GmbH, Germany.

Published: 18 May 2015

doi:10.1186/2197-7364-2-S1-A46

Cite this article as: Vaz et al.: Assessment of the quality of brain regions and neuroimaging metrics as biomarkers of Alzheimer's Disease. *EJNMMI Physics* 2015 2(Suppl 1):A46.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
